

Submitted by: William Rilling, MD, FSIR, Professor of Radiology & Surgery, Medical College of Wisconsin
Sean Tutton, MD, FSIR, Professor of Radiology, Surgery, & Palliative Medicine, Medical College of Wisconsin
Kristofer Schramm, MD, Assistant Professor of Interventional Radiology, University of Colorado

Organization:
Society of Interventional Oncology
2025 M St NW #800, Washington, DC 20036
[Phone: \(202\) 367-1164](tel:(202)367-1164)
[Email: TGreene@sio-central.org](mailto:TGreene@sio-central.org)

July 13, 2020
NCCN Guidelines Panel: Palliative Care

On behalf of the Society of Interventional Oncology, we respectfully request the NCCN Palliative Care Guideline panel review the enclosed data for inclusion in the NCCN Palliative Care Guidelines: inclusion of the recommendation for Interventional Radiologist (IR) consultation along with pain management and palliative care for all oncology patients.

Specific Change 1: Category 2A – Under unacceptable outcomes add consultation with an interventional radiologist along with pain management or palliative care specialist.

Rationale: Interventional Radiologists have a significant role in diagnosis and management of many cancer patients via both therapeutic and diagnostic procedures.

The increasing involvement of the interventional radiologist in care of the cancer patient has led to many novel treatment regimens as well as palliation strategies. , Palliative procedures such as vertebroplasty, fluid management (ex. pleural, abdominal and venting drainage catheter placement), and pain control procedures (ex. Celiac Plexus, hypogastric, intercostal, pudendal blocks) are underutilized but highly beneficial procedures that are provided by interventional radiologists. Percutaneous image guided ablation (both radiofrequency ablation and cryoablation) have been shown to provide excellent pain relief for skeletal metastases.

Change references

1. Callstrom MR, Dupuy DE, Solomon SB, Beres RA, Littrup PJ, Davis KW, et al. Percutaneous image-guided cryoablation of painful metastases involving bone: multicenter trial. *Cancer*. 2013;119(5):1033-41.
2. Position Statement on Percutaneous Vertebral Augmentation: A Consensus Statement Developed by the Society of Interventional Radiology (SIR), American Association of Neurological Surgeons (AANS) and the Congress of Neurological Surgeons (CNS), American College of Radiology (ACR), American Society of Neuroradiology (ASNR), American

William Rilling, MD
President
*Medical College
of Wisconsin*

Matthew
Callstrom, MD, PhD
President-Elect
Mayo Clinic

Stephen Solomon, MD
Immediate Past
President
*Memorial Sloan-
Kettering Cancer Center*

Muneed Ahmed, MD
Treasurer
*Beth Israel Deaconess
Medical Center/Harvard*

Jack Jennings, MD, PhD
*Washington University
School of Medicine*

Alexis Kelekis, MD
*Attikon University
Hospital Athens*

Kevin Kim, MD
Yale Cancer Center

Martijn Meijerink,
MD, PhD
Amsterdam UMC

Uei Pua, MBBS
Tan Tock Seng Hospital

Constantinos T.
Sofocleous, MD, PhD
*Memorial Sloan-
Kettering Cancer Center*

Ex-Officio
Michael C. Soulen, MD
*Hospital of the University
of Pennsylvania*

Executive Director
Jena Eberly Stack

Society of Spine Radiology (ASSR), Canadian Interventional Radiology Association (CIRA), and the Society of NeuroInterventional Surgery (SNIS)":

3. Mileo-Ferrer, Laura, Blanco Ana Isabel Luque, Gonzalez-Barbotoe Jesus. Efficacy of Cryoablation to Control Cancer Pain: A Systematic Review. Pain Pract 2018 Nov;18(8):1083-1098. doi: 10.1111/papr.12707. Epub 2018 Jun 7.

SIO also respectfully suggests that NCCN consider having an interventional radiologist participate as a panel member to add further multidisciplinary input and perspective on future versions of these guidelines.

Sincerely,
William Rilling, MD FSIR
Sean Tutton, MD, FSIR
Kristofer Schramm, MD, FSIR